

**Western Watersheds Project, Inc.**  
P.O. Box 280  
Mendon, Utah 84325  
435-881-1232 • [utah@westernwatersheds.org](mailto:utah@westernwatersheds.org)



March 23, 2004

Vernal BLM Field Office  
c/o SWCA Environmental Consultants  
230 South 500 East, Suite 380  
Salt Lake City, Utah 84102

Att: Ms. Deb Reber, Project Manager

Re: Vernal BLM Resource Management Plan DEIS (RMP/EIS)

Dear Ms. Reber:

Western Watersheds Project (WWP) is a 501c3 non-profit membership conservation organization with offices in California, Idaho, Montana, Utah and Wyoming. WWP, on behalf of all its members, has been working for over a decade to beneficially influence the management of BLM administered lands in the western United States. WWP has a long history of close involvement as an interested public on numerous grazing allotments administered by the BLM in Idaho, Nevada, Utah, Wyoming, eastern Oregon and southwest Montana.

WWP is submitting these comments and analysis for incorporation into the analysis of Range (Livestock Grazing) issues in the RMP/EIS. We support sustainable multiple use without loss of the potential productivity and diversity of these lands. In accordance with BLM's multiple use mandate, you must include consideration of the intrinsic values of the native biodiversity of these lands and protect those values for the long-term benefit of the American people as described in 43 CFR 1601.0-5(f).

WWP is knowledgeable of literally hundreds of grazing allotments which are failing the most minimal of environmental health criteria because of livestock grazing on BLM administered lands. The evidence we provide in these comments makes the case that these lands continue to be severely overstocked with livestock. BLM's own data and current management shows this to be the case. The best quantitative and peer-reviewed range science shows that continued emphasis on structural facilities erroneously called "range improvements" is a flawed strategy. BLM typically proposes these projects rather than making the difficult decision to adjust livestock numbers and seasons to be within the current capacity of the land. Without addressing this issue, the productivity and diversity of the land will continue to fall with the result that the land, the public interest and livestock producers will suffer over the long term.

As part of these comments, we refer to voluminous scientific literature and reports that BLM must consider in its analysis. Your review and analysis of these documents and inclusion of them in the RMP/EIS is necessary to provide a "balanced" approach to the issues. Their inclusion is also essential for BLM to comply with NEPA's

mandate to take a “hard look” at science and do a thorough and integrated analysis of all disciplines. WWP will provide copies of these publications to you for our costs of reproduction and postage.

Our comments are organized by sections including:

1. Introduction
2. What Does Range Science Tell Us?
3. Management Recommendations for Uplands
4. Management Recommendations for Riparian Areas
5. References

## **1.0 Introduction**

WWP is concerned that, in the past, BLM has not included consideration of the best available science in its environmental analyses. BLM does not follow the well known and science-based principles of range management in managing livestock grazing. The three pertinent federal statutes (FLPMA, PRIA and the Taylor Grazing Act) require that BLM administer these lands in the long-term interests of the American people and not a handful of stockmen, who are permittees, on the public lands.

Livestock permittees are a small minority of livestock producers in the eleven western states and are insignificant in their numbers or their economic contribution to the States, their local and regional economies. Their numbers and contribution pale in comparison to the natural values of our public lands. Dr. Thomas Power, Chairman of the University of Montana’s Economics Department, in Wuerthner and Matteson (2002) points out the minimal economic contribution of federal public lands livestock grazing to local, state and regional economies in the West, including Utah. That reference can be found on-line at:

[http://www.publiclandsranching.org/htmlres/PDF/wr\\_TAKING\\_STOCK.pdf](http://www.publiclandsranching.org/htmlres/PDF/wr_TAKING_STOCK.pdf)

Dr. Power also points out that the majority of public lands livestock producers depend on non-agricultural sectors of these local, state and regional economies for employment, not livestock production. It is not in the public’s interest to blindly continue livestock grazing at unsustainable stocking levels in order to provide a short-term benefit to this small minority, while ignoring the values displaced by livestock grazing.

In its environmental analysis and subsequent land use plan, BLM must include detailed consideration of the information and data provided in several recent publications and the references cited herein before making its decision. The following three books provide volumes of meaningful assessments of the unfortunate current condition of public lands and the failing economic and social realities of public lands ranching. These are:

- *Welfare Ranching, The Subsidized Destruction of the American West* (Wuerthner and Matteson 2002)

- *The Western Range Revisited: Removing Livestock from Public Lands to Conserve Native Biodiversity* (Donahue 1999)
- *Waste of the West* (Jacobs 1991)

In addition, BLM must consider in detail the publications referenced in these comments and also National Research Council (2002), *Riparian Areas: Functions and Strategies for Management*. All of these publications are clear in describing the flaws in current methods of livestock management on public lands and should form a core of its analysis.

In addition, the BLM's *Rangeland Reform '94 DEIS and Executive Summary* (RRDEIS, BLM 1995) reported that riparian areas "*have continued to decline and are considered to be in their worst condition in history*"; livestock grazing is identified as the chief cause. Indeed, some riparian areas have literally been destroyed; that is, they no longer exist or have any potential for restoration.

In its recent DEIS to revise its grazing regulations, BLM bypassed consideration of best available science. It pretended that current grazing management is fine if interested publics and environmental organizations would just quit holding up BLM and Permittee proposals by creating an administrative paperwork burden (BLM 2003). BLM asserted that greater cooperation with stockmen would somehow result in improvement by merely "tweaking" stocking rates, but then went on to state that this seldom happens, that changes in grazing really only amount to changes in season or location of use and that "*Changes in active grazing use in excess of 10% are infrequent.*"

In its RRDEIS, BLM provided definitions describing the status of upland plant communities. These were:

- Potential Natural Community (PNC) = existing vegetation is between 75 – 100% of the sites potential natural plant community.
- Late Seral Community = existing vegetation is between 50 – 74% of the sites' potential natural plant community
- Mid Seral Community = existing vegetation is between 25 – 49% of the sites' potential natural community
- Early Seral Community = existing vegetation is between 0 – 24% of the sites' potential natural community.

Table 1 indicates that management of uplands since the passage of the last Rangeland Reform regulation in 1995 has not resulted in improvement of upland condition. In fact, by BLM's own definitions, condition has declined. It also shows that productivity of those lands is greatly below potential with 63% of its lands below 49% of potential. The data also show that BLM's management has failed in the intervening 10 years to take meaningful actions to improve conditions.

In spite of the evidence of widespread loss of plant productivity and ground cover, accelerated erosion and BLM's own documentation of rapid declines in species such as sage grouse, BLM routinely chooses not to address livestock impacts in any scientific or sustainable fashion. Instead, BLM proposes more water developments and grazing

systems. This ignores that in the 1960's, BLM began a massive program of developing water, putting streams and springs into pipelines, seeding with crested wheatgrass, building fences, engaging in rotation grazing, and spending millions of dollars to "even out livestock distribution".

**Table 1. Comparison in BLM Upland Condition Between RRDEIS (1995) and Current Condition (BLM 2003)**

<b>Community Status</b>	<b>RRDEIS</b>	<b>Current DEIS</b>	<b>Change '94 to date</b>
PNC	4%	6%	+2%
Late Seral	34%	31%	-3%
Mid Seral	40%	34%	-6%
Early Seral	15%	12%	-3%
Unclassified	7%	17%	+10%

An early example of this, among others, was in BLM's Vale District, where millions of dollars were spent on crested wheatgrass seedings and structural range improvements. Today, across BLM lands in the west, many of these systems have fallen into disrepair, the land has failed to recover and we are faced with more and more proposals to install grazing systems, water developments, plow and seed – not reduce livestock numbers. This is in spite of the fact that long-term studies, including those from the Vale District have shown that stocking rate is the critical variable, not grazing systems. These are cited in a later section.

This is all in the context of BLM's failure to scientifically and accurately determine those lands which are capable and suitable for livestock grazing. We must add to this the further failure of BLM to accurately and quantitatively determine how much forage (i.e. forage capacity) is currently available. On top of this, there is the failure of BLM to properly allocate that forage to watershed and stream protection, wildlife habitat and food, then to livestock if available. Then there is the failure to provide for long-term rest to facilitate recovery. Finally, we must add the unwillingness of permittees to use peer-reviewed range science principles for management, rely on "snake-oil" solutions such as time-controlled grazing and their strong opposition to the most minimal standards of performance. These failures by BLM and livestock permittees have prevented the recovery of damaged ecosystems in order that they might sustain use as envisioned in the Taylor Grazing Act and FLPMA.

Instead, BLM continues to use the take "half/leave half" principle for livestock use. In its permits and analyses, BLM typically includes livestock use levels of 50% of forage as proper. They do this without providing any scientific foundation for this claim, nor do they adequately monitor this use or use it to manage livestock. The following paragraphs provide a summary of the relevant range science regarding utilization levels, plant growth and productivity, effects of precipitation regime, capability and suitability, capacity determinations, stocking rates and range economics. These principles are well founded in the range science literature.

## 2.0 What Does Range Science Tell Us?

**2.1 Plant Growth and Precipitation.** In order to understand the implications of grazing livestock at these heavy forage utilization levels in arid regions, it is important to understand the precipitation regime and its relationship to forage production. Holechek et al (2001) point out what we all understand at the most basic level. That is, precipitation is the single most important factor determining the type and amount of vegetation in a particular area. In the 11 Western States, 80% of the area receives less than 500 mm (19.6 inches) of annual average precipitation. Further, this precipitation is subject to great year-to-year variation.

Three locations representative of precipitation patterns for Utah BLM lands are used for illustration. Long-term precipitation records were analyzed for stations at Escalante, Kanab and Vernal. The effects of annual variations in precipitation on plant community production are important to understand in establishing livestock stocking rates and management. Table 2 provides a summary of annual precipitation statistics for these locations. Data was obtained from the Western Regional Climate Center database which can be found on line at <http://www.wrcc.dri.edu/index.html> . Figure 1 provides plots of annual precipitation and percent occurrence of below average and drought years for these locations.

The analysis uses the Standard Precipitation Index (SPI) developed by McKee et al (1993) which considers drought, or extremely dry conditions, as years with 2" less than average precipitation.

**Table 2. Summary of Precipitation Statistics for Three Utah Locations**

Description	Escalante	Kanab	Vernal
Period of Record	1909 – 2001	1903 – 2002	1928 – 2003
Years of Record	65	86	70
Average, inches	10.9	13.4	8.2
Range, inches	5.5 – 21.7	5.4 – 26.6	2.8 – 14.8
Year below average	39	45	33
Percent years below average	60%	52.3	47.1
No drought years based on SPI	18	31	13
Percent drought years based on SPI	27.7	36	18.6

Figure 2 shows the monthly distribution of precipitation at these locations. The differences in overall precipitation amounts are reflected in the relative magnitudes of monthly precipitation at each of the locations, with Vernal being the driest. Escalante and Kanab have generally drier spring and early summer periods, while Vernal experiences an increase in precipitation during spring, then a decline. All locations

have increased precipitation during late summer and early fall. Winters are generally characterized by consistent precipitation from month to month.

These periods of precipitation vary in their effects on the plant communities. Increased precipitation during the spring-summer-fall months may not be effective due to the higher temperatures occurring then. Typically, the fall-winter period is the period of greatest increase in soil moisture due to lower temperatures and lower evapotranspiration. Precipitation effectiveness during the warmer spring – fall periods varies with the storm intensity and soil condition. Storms must be of high enough intensity to promote recharge of the soil profile into the root zone to be effective for plant growth. Generally, this is greater than 0.6 inches in desert shrub types, although very high intensity storms may not be effective due to rainfall rates in excess of infiltration that result in overland runoff and flash events.

Spring plant growth in these arid areas depends on the amount of moisture received and retained during the fall-winter period. Relatively dry summers may allow little regrowth and by the time fall comes, temperatures may be low and growth limited. Trampled and compacted soils exacerbate this effect (Blaisdale and Holmgren 1984). Some desert shrubs such as *Artemisia* sp. with both shallow and deep root systems can take advantage of both shallow and deep soil moisture (West 1983).

Annual production of available forage at the Desert Experimental Range in western Utah was highly correlated with total annual precipitation, showing an 800% variation in forage production between the driest and wettest years (Hutchings and Stewart, 1953). Scientists developing quantitative ecosystem relationships for the Prototype Oil Shale Program managed by BLM in Utah's Uinta Basin found that annual sagebrush stem leader growth used as an index of production had a high correlation with winter precipitation (October – March) and that spring annual plant biomass was correlated with spring precipitation (ERI 1984; WRSOC 1984). These studies were carried out under the supervision of BLM's Vernal Field Office and the data and reports that came from that detailed characterization of the environment are available for inclusion in preparation of the RMP/EIS.

Analysis of twenty years of data for perennial grass production and annual precipitation for a study area at the Chihuahuan Desert Rangeland Research Center in New Mexico showed a high correlation (Holechek et al, 2001). A graph of this data is shown in Figure 3. Annual perennial grass production varied between 6 and 750 lbs per acre, corresponding to the second-lowest and highest precipitation years. The linear regression plot of the same data is provided in Figure 4. Results of long term studies of crested wheatgrass production from experimental plots on BLM land at Malta, Idaho showed that crested wheatgrass production was most closely related to May-June precipitation (Sharp et al, 1992). They found that annual production of crested wheatgrass during 35 years averaged about 500 pounds/acre and ranged between 130 and 1090 pounds/acre depending on precipitation (Figure 5). These relationships demonstrate that this is a predictable phenomenon that should be taken into account in setting livestock grazing seasons, stocking rates and management on an annual basis as well as over the longer term. A typical NRCS soil survey (USDA 1980) for these arid areas shows that total production of potential plant communities varies by about 300% between favorable and unfavorable years. This wide range in production between dry and wet years is typical in the arid regions of the West.







Figure 1. Annual precipitation patterns in Escalante, Kanab and Vernal, Utah








Figure 2. Monthly precipitation patterns in Escalante, Kanab and Vernal, Utah


**Figure 3. Annual Precipitation vs Forage Production**



**Figure 4. Forage Production vs Precipitation**



**Figure 5. Crested Wheatgrass Production at Malta, Idaho**



**2.2 Grazing Intensity and Effects on Productivity.** Much of the current research and analysis of livestock grazing management, plant productivity and economics has come out of the Department of Animal and Range Sciences at New Mexico State University. This work has been presented in a series of textbooks and papers in the range science literature. These references provide analyses of the interactions of livestock stocking rates, plant productivity and economics based on a set of long term grazing management studies from native rangeland types. They provide recommendations for determining livestock grazing intensity to maintain vegetative productivity and economic stability, while taking into account the effects of inherent variation in precipitation in desert ecosystems.

The effects of different livestock grazing intensities on forage plant production was studied in a ponderosa pine type in Colorado as early as the 1940's (Schwan et al, 1949). This study showed that forage consumption at a rate of 57% produced an average of twice as much forage as a rate of 71%. An area left ungrazed by livestock for 7 years produced three times as much forage as the 71% use area. The authors concluded that, as grazing use increased, forage production decreased. During that same period, Dyksterhuis (1949), in a classic paper on the use of quantitative ecology in range management, presented examples of how stocking rates must be adjusted based on precipitation and range condition, which included a rating based on departure from the potential plant community. NRCS (USDA, 1982) considers proper grazing management as that management that sustains the potential plant community.

The effects of conservative (30 – 35%) use vs. heavy (60 – 65%) grazing use on grasses and forbs by cattle was determined in a New Mexico study (Galt et al, 1999). Both of these pastures had experienced conservative use for over 10 years. In 1997, one pasture was changed to heavy use. Quantitative measurements at key locations in both pastures in the following year, while being rested, provided the results shown in Table 3.

**Table 3. Standing Crop of Grasses and Forbs from Galt et al (1999)**

<b>Location/Forage Component</b>	<b>Spur Pasture Heavy Stocking Rate Pounds/acre</b>	<b>Deep Lake Pasture Conservative Stocking Rate Pounds/acre</b>
Perennial Grasses	352	824
Forbs	256	436
Total Forage	608	1260

This study showed that heavy stocking rates resulted in serious declines in productivity in the succeeding year. Perennial grass production was reduced by 57% and forbs by 41% in the heavily grazed pasture compared to the conservatively grazed pasture. The authors cited a number of other studies in arid environments that showed heavy stocking rates were accompanied by decreases in forage production when compared to conservative use. After drought, the ability of forage plants to recover was directly related to the standing crop levels maintained during the dry period. The studies cited showed that grazing during different seasons was less important than grazing intensity.

Five long-term stocking rate studies from three different locations in Arizona, New Mexico and Utah documented similar patterns (Holechek et al 1999a). In the Desert Experimental Range in Utah, a 13-year study with moderate (35%) and heavy (60%) use by sheep resulted in annual forage production of 198 lbs/acre and 72 lbs/acre. The authors recommended 25 – 30% use of all forage species. A 10-year study at the Santa Rita Range in Arizona demonstrated that perennial grass cover and yield showed an inverse relationship to grazing intensity, while burroweed, an undesirable species, increased with increasing forage use. The authors recommended a 40% use level. A 37-year study at the Jornada Experimental range in New Mexico involving conservative (33%) and moderate (45%) use showed that the lower grazing intensity resulted in greater black grama (perennial grass) cover. Lowland areas with high clay content and periodic flooding grazed at moderate intensity had higher cover of Tobosa, a perennial grass, than heavily grazed areas. They recommended 30% be used as a stocking intensity with no more than 40% removed in any year. A 10-year study at the Chihuahuan Desert Rangeland Research Center looked at four grazing intensities of 25%, 35%, 50% and 60%. Light (25%) and moderate (35%) use produced 70% more forage than 50% use and more than double that achieved at 60% use. Here, the author recommended conservative stocking at 30 – 35%.

Hutchings and Stewart (1953), suggested that 25 – 30 % use of all forage species by livestock was proper. They recommended this level because routinely stocking at capacity will result in overgrazing in half the years and necessitate heavy use of supplemental feed. Even with this system, they recognized that complete destocking would be needed in 2 or 3 out of ten years. Holechek et al (1999a) concluded that the research is remarkably consistent in showing that conservative grazing at 30 – 35% use of forage will give higher livestock productivity and financial returns than stocking at grazing capacity. They also recognized that consumption by rodents and other wildlife must be taken into account as part of this utilization. Otherwise, rangeland productivity would suffer even at these levels of use. Galt et al (2000) recommended levels of 25% utilization for livestock and 25% for wildlife with 50% remaining for watershed protection. In none of these cases have the scientists recommended 50% utilization by livestock and they are clear that even at the lower use levels recommended, wildlife use is included.

BLM has never even looked into the issue of the relationship between its typical riparian greenline stubble height standard of 3 – 4”, usually applied to Nebraska sedge and other sedge species, and the effects of applying this standard on adjacent riparian plant community productivity. Carter (1998) showed that before greenline N. sedge had been reduced in height to the standard of 6”, riparian grasses had been reduced to 2” stubble height and 89% of stream banks had been trampled, compacted or were actively eroding into the stream. Monitoring data from the Wasatch-Cache National Forest (USDA 1993) included the observation that a Nebraska sedge stubble height of 7.6” corresponded to an estimated 70% use of riparian grasses. Lile et al (2003), compared clipping of N. sedge to stubble heights of 2” and 4” during early season, late season and multiple clippings to both heights. Late season use or 2” stubble height did not allow recovery. Only the 4” early season use achieved the 4” criteria, but did not regrow to meet the 6” criteria by the end of the growing season. This shows that season-long or late season use does not allow sufficient time for re-growth and that 3-4” stubble heights are not effective in protecting riparian vegetation.

Schulz and Leininger (1990) studied long-term riparian exclosures compared to areas that continued to be grazed. They found that, after 30 years, willow canopy cover was 8.5 times greater in livestock exclosures than in adjacent grazed riparian areas. Grasses were 4 to 6 times greater in cover within the exclosure than outside. Mean peak standing crop of grasses within the exclosure was 2,410 Kg/Ha, while outside in caged plots, mean peak standing crop was 1,217 Kg/Ha.

Often cited, Franklin Crider's study on root growth stoppage from plant top removal provided quantitative measurements of plant re-growth under different amounts of removal (Crider 1955). Three mid-west perennial grasses were grown from seed in pots under ideal conditions of watering and fertilization. After sixty days of growth, these potted grasses were clipped once at intervals from 10% to 90% of the above ground biomass. Repeat clippings of the potted grasses were made every two days to return the plants to the same height as the original clipped percent. The experiment lasted thirty three days at which time root growth of controls became inhibited by the size of the pot. Crider concluded that under these ideal growing conditions, if these species of grasses had 40% or less of their aboveground biomass clipped either once or many times, then the net root mass was the same or more at the end of the experiment. This was used to make the assumption that grazing during the entire growing season at 40% or less would sustain plants from one season to the next. This same study has been used to justify the 50% or "take half/leave half" proposition that range managers have used for decades. Clearly, the long-term range studies cited here show that under actual field conditions, these use levels are excessive and light grazing (25%) is most equitable to BLM's mandate for sustainable use.

**2.3 Forage Needs Other Than Livestock.** Let's assume for the moment that Crider's finding (40% removal in a growing season) applies to the very arid lands found in the Vernal resource area. If Crider's results apply, then the 40% removed needs to be shared with all the animals that require forage. Some is needed for wild grazers (insects, nematodes, mammals), some for the generation of litter and nutrient cycling. A certain amount of the plant needs to remain to ensure regeneration of the plant and to ensure plant community structure is sufficient for wildlife habitat needs and soil erosion protection. A certain proportion of plants must remain ungrazed to allow for seed and flower production to ensure propagation of the species and provide food for pollinators and granivores, an often overlooked requirement.

Existing research helps answer the question of how much of the annual plant growth is required for wild grazers. For a detailed review of this research we recommend that you download Catlin et al (2003) from:

<http://rangenet.org/directory/jonesa/sulrprec/index.html>

Based on a number of studies of mammal consumption of forage, some annual forage consumption needs for mammals where mammal populations are at or near their potential are provided in Table 4:

**Table 4. Forage Needs of Mammals**

<b>Mammal</b>	<b>Forage Needs kg/hectare</b>
Deer	12.7
Rabbits	72.7
Rodents	142.3
Total	227.6

Less is understood regarding the typical vegetation needs for insects. What is known is that insects play a significant role. For example, harvester ants are granivores with individual colony populations of tens of thousands of individuals. Even in the driest and hottest parts of North America, the total biomass of one species of harvester ant, *Messor (Veromessor) pergandei* has approximately the same total biomass as the total rodent population in the same area. One of many ecological roles harvester ants fulfill is dispersing plants as a result of accidental abandoning of seeds near the nest. The analysis here focuses on the typical forage needs of insects.

A number of studies have reported the percent of plant growth consumed by insects. In grassland and savannah, insect consumption of a single species varied from less than 1% to 19% with a mean of 3.5% (Wiegert and Peterson, 1983). Rangeland grasshoppers consume between 1- 3% in typical years (Mispagel 1978) and up to 99% in periods of super abundance (Nerny and Hamilton 1969). Hewitt and Onsager (1983) suggest that between 21-23% of available range forage is consumed by all grasshopper species in the western U.S. each year. This production is transferred to birds and mammals such as sage grouse and rodents as a food source, to the soil for nutrient cycling and to higher trophic levels such as raptors and other carnivores. It is important for ecosystem function and cannot be dismissed.

Nematodes provide critical soil nutrient cycling and structure functions. This complex array of species normally has a higher diversity and biomass than other rangeland flora and fauna combined. Some nematodes consume plants and should be considered as a component in the consumption of rangeland plants. Ingham and Detling (1984) estimated that nematodes consumed between 6 and 13 % of below ground net productivity in the mixed grass prairie. Other studies by Scott et al (1979) and Anderson (1987) found that root feeding arthropods and nematodes consume between 7-26% of the net productivity in normal years. For the full citations to these studies please visit the paper at the web site shown above.

Holechek et al (2001) provide equivalents in forage consumption between livestock and wildlife. They are summarized in Table 5. Studies in the Vernal resource area during the 1970's and 1980's generated many years of quantitative data describing the abundance of many species of animal, bird and insect in the different vegetation types occurring there. This information is available in the Vernal Field Office for review and analysis. Some of it is provided in the (ERI 1984; WRSOC 1984) reports referenced. Final determinations of livestock stocking rates and utilization criteria for livestock must take this information into account, using the current vegetation production for the area under consideration.

**Table 4. Animal Equivalents from Holechek et al (2001)**

<b>Animal</b>	<b>Animal Unit Equivalents</b>
Mature cow	1.0
Yearling cow	0.75
Domestic sheep	0.15
Horse	1.8
Bison	1.8
Elk	0.7
Moose	1.2
Bighorn sheep	0.18
Mule deer	0.15
Pronghorn	0.12

**2.4 What is an AUM?** Various estimates have been used in the past to define the amount of forage consumed by livestock (AUM). BLM often uses the value of 800 lbs/AUM. This section discusses some of that variability and proposes a standard forage amount for livestock consumption. A check of USDA market statistics at:

[http://www.ams.usda.gov/mnreports/lm\\_ct166.txt](http://www.ams.usda.gov/mnreports/lm_ct166.txt)

shows that during 2003, auction weights of mature steers averaged 1268 lbs and mature heifers averaged 1157 pounds. Recent auction weights for mature cows have seen ranges up to 1400 pounds. Ray et al (2004) give a weaning weight of 480 pounds for calves. Holechek et al (2001) state that daily dry matter consumption averages 2% of body weight per day. They then calculate that a 1000 pound cow will consume 20.0 lbs dry forage/day and a 750 pound yearling will consume 15 lbs dry forage/day, for a total of 35 lbs/day. The 1969 AMP for the North Rich Allotment in Rich County and Cache Counties (Wasatch-Cache NF) used 30 lbs/day for a cow calf pair. In its FEIS for the Land and Resource Management Plan for the Curlew National Grassland (USDA 2002), the Caribou-Targhee National Forest used 34 pounds per day for a cow/calf pair. There is little doubt that cattle weights have increased over time and a value that is accurate under current conditions must be applied. In fact, Anderson et al (ca 2000) calculated a 35% increase in dressed weights per animal between 1975 and 1995. Their analysis can be found at:

<http://agecon.uwyo.edu/RiskMgt/marketrisk/TheCattleCycle.pdf>

Using the average auction weight for heifers (1157 lbs), the weaning weight for calves (480 lbs) and the 2% consumption rate from Holechek et al (2001) results in forage dry weight consumption of:

$$(1157 + 480) \times 0.02 \times 30 \text{ days} = 982 \text{ lbs/month, or } 32.7 \text{ lb/day}$$

Even this will likely be an underestimate since calves graze all summer and are probably closer to the yearling weight given in Holechek et al (2001) and mature cows can be a good deal larger than the heifer weight used here. We propose that BLM

establish a forage consumption allocation for a cow/calf pair of 1,000 lbs/month = 1 AUM.

**2.5 Grazing Systems.** In a review paper that considered grazing systems, grazing intensity and season of use, Holechek et al (1998) determined that, *“financial returns from livestock production, trend in ecological condition, forage production, watershed status and soil stability are all closely associated with grazing intensity.”* They found that grazing systems such as rest-rotation had limited or no benefit in arid systems. Citing long-term studies in Arizona, they documented that after 12 years of rest-rotation management compared to continuous grazing, neither forage plant densities nor forage plant production differed between the treatments. Grazing intensity employed was 30 – 35% use with occasional high use of 50% or more. *“Rest and deferment were not sufficient to overcome the effects of periodic heavy use on primary forage plants when rest-rotation grazing was applied on big sagebrush range in northern Nevada.”* In an Arizona study comparing winter-spring grazing with summer-fall rest to continuous grazing, the rotation scheme was inferior to the year-long system from the standpoint of perennial grass density and production. Perennial grass production was closely associated with the degree of use and was highest where grazing use was lowest. In a Vale, Oregon study, lasting over 20 years at moderate grazing intensity, rotational grazing showed no advantage over season-long grazing in improving range condition or forage production. *“The key factor in range improvement appeared to be the reductions in grazing intensities that were applied when the project was initiated..”*

A review of the “classic” range studies, which are the long-term stocking rate and grazing system studies that provide the scientific foundation for modern range management again showed that light use is closer to sustainable use, while heavy use is not (Holechek et al 1999a). Definitions of “heavy”, “moderate” and “light” grazing developed in 1961 were cited. Heavy grazing was defined as the degree of forage utilization that does not allow desirable forage species to maintain themselves. Moderate grazing was defined as the level at which palatable species can maintain themselves. Light grazing was defined as the degree of utilization at which palatable species are able to maximize their herbage producing ability. However, it is clear that using even “moderate” grazing in depleted areas will not allow them to recover.

When averaged across all the long-term studies for all regions, heavy grazing was 57% use of primary forage species, moderate use was 43% and light use was 32%. In arid regions, the research showed that moderate grazing use was 35 – 45%. When the average forage production change over time was compared with use, heavy stocking resulted in a 20% decline in production, moderate use experienced no change and light use resulted in an 8% increase. During drought, moderately stocked pastures produced 20% more forage than heavily stocked pastures, light grazing produced 49% more forage than heavy and 24% more than moderate stocking levels. Heavy stocking resulted in a downward trend and light stocking an upward trend in ecological condition. Moderate stocking showed a slight, but not significant increase in condition, resulting in depleted ranges being maintained in depleted condition.

Table 6 provides summary statistics from that paper. It must be remembered that these comparisons are to prior heavy use, not to ungrazed lands. It is apparent from these studies that “moderate” use levels will not allow significant recovery of severely

depleted range. In fact, in studies of long-term rest at Idaho National Engineering Laboratory, the recovery rate of grasses in sagebrush communities was slow, progressing from 0.28% to 5.8% over 25 years (Anderson and Holte, 1981 and Anderson and Inouye, 2001). It is clear from these examples that native plant communities in heavily depleted sites will require decades to recover in the absence of livestock, while their ability to recover in the presence of livestock at any level of use has not been demonstrated.

Relying on additional water developments, fences and grazing systems will not alleviate the problem. The use of range improvements and rotation systems is not sufficient to correct over-stocking. Results from 18 western grazing system studies by Van Poolen et al (1979) found that adjustment of livestock numbers, or stocking intensity was more important than implementing grazing systems to improve herbage production. Holechek et al (1999a) recognized that “*various rotation grazing systems cannot overcome the rangeland deterioration associated with chronic overstocking.*” Holechek et al (2000) also showed that the various claims made by advocates of short-duration or time-controlled grazing were false.

A comprehensive discussion of rest-rotation is found in Clary and Webster’s General Technical Report titled “*Managing Grazing of Riparian Areas in the Intermountain Region.*” (Clary and Webster 1989). They summarized a dozen studies showing significant increases in forage production occurred with decreased intensities of grazing. The article described the improvements found in reducing grazing from heavy, to moderate and then to light grazing. Grazing with utilization above 50% was described as heavy, moderate was 30 - 50% and <25-30% was called light grazing in most of these studies.

**Table 5. Summary of Data from 25 Classic Grazing Studies (Holechek et al 1999a)**

<b>Description</b>	<b>Heavy</b>	<b>Moderate</b>	<b>Light</b>
Average Forage Use %	57	43	32
Average Forage Production lb/acre	1,175	1,473	1,597
Drought Years Production lb/acre	820	986	1,219
Average Calf Crop %	72	79	82
Average Lamb Crop %	78	82	87
Calf Weaning Weight lbs	381	415	431
Lamb Weaning Weight lbs	57	63	--
Gain per Steer lbs	158	203	227
Steer/calf Gain per Day lbs	1.83	2.15	2.3
Steer/calf Gain per Acre lbs	40.0	33.8	22.4
Lamb Gain per Acre lbs	26.0	20.4	13.8
Net Returns per Animal \$	38.06	51.57	58.89
Net Returns per Acre \$	1.29	2.61	2.37

Clary and Webster's study concluded that *"managers should place more emphasis on proper stocking intensity and less on grazing system implementation. The concentrated use of grazing pastures is not compensated for during rest years if grazing use is heavy. In summary, although grazing systems have great intuitive appeal, they are apparently of less consequence than once thought. In fact as long as good management is practiced so that there is control of livestock distribution and grazing intensity, the specific grazing system employed may not be significant."*

Holechek et al (2001) have indicated that, depending on topography, areas of severe degradation, or *"sacrifice areas"* occur around water sources including water developments. These can extend from 1 mile to several miles from these sources and out further if stocking rates are too high. Based on this, a single water development can result in an area of soil compaction, erosion and severe loss of ground cover and vegetation for thousands of acres. They also indicate that installing water developments in locations that have had limited access to livestock in the past may increase ecological damage to areas that are important refuges for relict plant communities and wildlife that have not been displaced by livestock.

Appendix B in Catlin et al (2003) provides a review of grazing systems. It summarizes more than 40 studies concerning rest rotation and related issues. Short term rest (one year) as typically applied does not lead to significant improvements of deteriorated ranges. Clearly, the long-term range studies we have cited show that it is stocking rate, not water developments, or grazing systems that are most important in maintaining or improving rangeland productivity. It is critical that BLM take this information into account in determining livestock stocking intensities and evaluating the efficacy of added water developments or grazing systems.

**2.6 Economic Considerations.** The studies we have cited show that light stocking results in greater forage production and improvement in range condition when compared to both heavy and moderate use. Moderate and light use also provided greater financial returns than those obtained with heavy use. Because these financial figures included data from humid areas, a separate analysis taking into account the necessity for destocking during drought in arid regions showed that conservative stocking (35% use) would provide the highest long-term financial returns on semi-desert rangelands in Arizona.

Economic analyses in (Holechek et al, 1999b) show that conservative stocking rates yield better returns. For example, in the sheep experiment at the Desert Experimental Range in Utah, the lower stocking rate (35% use) yielded a financial return of \$0.39/acre compared to \$0.14/acre for the higher stocking rate (60% use). A modeling study that evaluated 29 years of financial returns for a cow-calf operation revealed that a relatively constant stocking rate of 35% use was considered the best approach.

Winder et al (2000) reported on comparisons of stocking rates and financial returns using 30 or 40% of current years perennial grass growth. The 30% use level provided greater vegetation productivity and financial returns. After drought in 1994 through 1996, forage production on the pastures with the lower stocking rate (30% use) increased 71% compared to 35% increase on those with the moderate stocking rate

(40%). Economic returns were \$0.52/acre for the conservative use level and \$0.31/acre for the moderate use level. A combination of stubble heights, clippings and ocular measures were used to set annual stocking rates, termination of the grazing season, sale of cattle to balance numbers with capacity and destocking during drought. Under these criteria, all pastures were destocked in the summer of 1994 and the moderately stocked pasture was destocked in May, 1999. After livestock were removed due to drought, pastures were rested for two years, then afterwards stocked in late fall according to current year's forage production.

Results of seven years' research in New Mexico's Chihuahuan Desert to evaluate the relationship between range condition and financial returns showed similar relationships (Holechek et al 1996a). Condition was evaluated using the Dyksterhuis (1949) definitions based on departure from climax. This study showed a relationship between forage production and range condition. Higher condition range, or that nearer climax community plant composition, had higher production of forage and more preferred forage species than lower condition range. Excellent range condition provided over four times the financial return of fair condition range and 65% greater return than good condition range.

The reasons for this were the high costs of management and the energy lost by livestock in seeking forage in lower condition range. In a companion paper, livestock returns were compared to conventional investments such as bonds or stocks (Holechek et al, 1996b). This analysis showed that over-capitalization in infrastructure, coupled with over-stocking lead ranchers into a boom and bust cycle as climatic conditions change. In wet years, they added livestock, generally when prices were high then sold off their herds during dry, or bust periods when prices and productivity are low. The final analysis concluded that conservative stocking, minimal investment in range improvements and greater spacing of watering points reduced fixed costs and insulated the operation from the vagaries of precipitation and market forces.

In addition to the economics of livestock production, BLM is required to analyze other economic values and determine the values foregone by its proposed action. NEPA (40CFR1508.8) recognizes that the analysis of effects must include ecological, aesthetic, historic, cultural, economic, social, or health issues. BLM typically has clouded the distinction between "cultural" as applied to historical features such as buildings, artifacts, or paleo-resources with societal lifestyle issues. In particular, in its recent DEIS to revise the grazing regulations, BLM used the abstraction called "lifeways" . This distinction needs to be clearly drawn. It appears BLM is trying to draw in "lifestyle" or "lifeways" as some sort of cultural feature that is given protection by the cultural preservation laws and SHPOs. This is not the intent of these laws or NEPA. If it were, would not "lifestyles" or "lifeways" other than livestock production also deserve protection, consideration and analysis?

NEPA, at 40 CFR 1501.2 (b) states that federal agencies must *"Identify environmental effects and values in adequate detail so they can be compared to economic and technical analysis."* 43 CFR 1601.0-5(f) defines Multiple Use as: *"Multiple use means the management of the public lands and their various resource values so that they are utilized in the combination that will best meet the present and future needs of the*

*American people; making the most judicious use of the lands for some or all of these resources or related services over areas large enough to provide sufficient latitude for periodic adjustments in use to conform to changing needs and conditions; the use of some lands for less than all of the resources; a combination of balanced and diverse resource uses that takes into account the long term needs of future generations for renewable and non-renewable resources, including, but not limited to, recreation, range, timber, minerals, watershed, wildlife and fish, and natural scenic, scientific and historical values; and harmonious and coordinated management of the various resources without permanent impairment of the productivity of the lands and the quality of the environment with consideration being given to the relative values of the resources and not necessarily to the combination of uses that will give the greatest economic return or the greatest unit output.”*

FLPMA requires BLM to manage the public lands for multiple use. This means that the agency must make “*the most judicious use of the land for some or all of [the various] resource values . . . [and to] use . . . some land for less than all of the resources . . . .*” 43 U.S.C. § 1702(c). This requires BLM to undertake a decision process “*to balance competing resource values to ensure that the public lands are managed in a manner that will best meet the present and future needs of the American people.*” 43 U.S.C. § 1702(c); Comb Wash at 101. BLM’s balancing of values must be reasoned and well-informed. Therefore, the agency must accumulate sufficient data and consider relevant rigorous science to determine what uses are appropriate in any given area.

Applied to the management of livestock grazing, the analysis must, on a site-specific level, weigh the benefits and harms of grazing to determine if BLM should allow this use in any given area. Moreover, should the agency conclude that livestock grazing is an appropriate use, BLM must consider multiple-use values in determining how that area should be grazed. National Wildlife Fed’n v. BLM, No. UT-06-91-1 (DOI, Office of Hearings and Appeals, Hearings Div.) (Dec. 20, 1993) at 25, *aff’d Comb Wash* (citing 43 U.S.C. §§ 1701(a)(8), 1702(c)). Therefore, in establishing grazing thresholds such as stocking rates and utilization levels, BLM is required to abide by “*FLPMA’s mandate [that it] protect the full spectrum of environmental, ecological, cultural, and recreational values.*” Id.

Other than asserting in various ways that continued livestock grazing at current levels provides for preservation of rural values and lifestyles, BLM generally does not provide any economic analysis of the costs and benefits of public lands livestock grazing and its contribution to local and regional economies. It does not analyze the values of uses foregone in favor of livestock grazing and its infrastructure. See the detailed and quantitative analysis by Dr. Power in (Wuerthner and Matteson 2002) that is referenced above.

Dr. Power shows that “*Livestock grazing on federal lands is generally unimportant to local economies and even less so to state and regional economies. In terms of income and numbers of jobs provided, the contribution of federal lands grazing is less than 0.1% across the West. Farm and ranch operations are increasingly reliant on nonfarm income sources to be financially feasible, while livestock grazing competes with other uses of public lands – such as clean water, recreation, wildlife habitat – that contribute to the ongoing vitality of western economies.*”

In his analysis of the economies of individual rural counties, Dr. Power showed that federal lands grazing does not contribute significantly to those economies across the

west. In fact, given the high percentage of ranching families that have jobs, either full or part time outside the ranch (60 – 70%), it is ranchers that depend on the other economic sectors for their ability to persist, not federal grazing. Dr. Power states, “It is not that towns depend on agriculture, but that agriculture increasingly depends on the vitality of urban and nonagricultural rural economies to provide the nonfarm income that keeps farm operations alive.”

Dr. Power states that claims about the relative importance of federal grazing to the economies of western states can be analyzed by answering these questions:

1. “What portion of the value produced by cattle and sheep operations is associated with feed used?
2. What portion of the feed for those cattle and sheep operations comes from grazing on federal lands?
3. What portion of the total agricultural activity involves raising cattle and sheep?
4. What part of the total economy is represented by agriculture.”

Souder (1997) analyzed the economics of livestock grazing in the Central Winter Ecosystem Management Area on the Kaibab Plateau using readily available economic data. His analysis showed the values of various activities in the study area generated values that are tabulated in Table 7.

**Table 7. Summary of Values Received Annually in the Central Winter EMA**

<b>Resource</b>	<b>Local Benefits</b>
Dispersed Recreation	\$6,400,000 (Locally and Regionally)
Hunting Mule Deer and Turkey	\$719,392 (Mule Deer = \$470,528)
Livestock Grazing	\$43,283
Fuelwood Gathering	\$45,492

BLM should include consideration and analysis of these sources in its RMP/DEIS and also provide more analysis of the economic benefits of wildlife through hunting, fishing and wildlife-watching associated recreation. These benefits are summarized in the Fish and Wildlife Service *2001 National Survey of Fishing, Hunting and Wildlife-Watching Associated Recreation* (DOI 2002). That survey showed that in Utah alone, expenditures for hunting, fishing and wildlife-associated recreation were \$1,143,408,000.00 in 2001

**2.6 Grazing Capability and Suitability Determinations.** Current range science recommendations include adjusting the stocking rate for livestock in order to account for distance from water and steepness of slope (Holechek et al, 2001). The Natural Resources Conservation Service has adopted these guidelines for slope adjustments (Galt et al, 2000). Their suggested reductions in grazing capacity for cattle with distance to water and increasing slope are provided in Table 8.

They note that on cold desert ranges of the U.S., snow reduces water availability problems in winter. Also, sheep do not require water every day and can use areas further than 2 miles from water. Sheep on New Mexico winter ranges used slopes of less than 45% with no adjustment necessary for slope, whereas slopes greater than 45% were hardly used. Regional criteria for the Intermountain Region of the Forest

Service designate lands with greater than 30% slope as not capable for cattle and greater than 45% slope as not capable for sheep. Other factors used by the Intermountain Region of the Forest Service for determining lands that are not capable include: current vegetation production less than 200 lb/acre, forested areas and areas with highly erodible soils (Blackwell 2001; USDA 2001).

**Table 7. Adjustments for Distance to Water and Slope for Cattle (Galt et al, 2000)**

<b>Distance from Water miles</b>	<b>Percent Reduction in Grazing Capacity</b>
0 - 1	0
1 - 2	50
>2	100
<i>Slope %</i>	
0 - 10	0
11 - 30	30
31 - 60	60
>60	100

Suitability determinations should be performed on those lands that are found capable for livestock to determine whether or not livestock grazing should be allowed. For example, important or critical fish and wildlife habitat, recreation areas, locations of sensitive populations, natural research areas, watershed protection areas among others should not be considered suitable and should be closed to livestock. These capability and suitability determinations are critical components in meeting the definitions and mandates of FLPMA, PRIA and NEPA regarding sustainability and multiple use. See also the discussion in the previous section regarding balancing of resource uses and values.

### **3.0 Management Recommendations for Uplands**

It is critical for BLM to incorporate the science we have cited in its RMP/EIS and in future planning and permit documents to ensure that best available science is applied on the ground. In this way, it may be possible to sustain livestock grazing on those lands that are capable and suitable. The following steps should be implemented in the RMP/EIS.

**3.1 Literature Review.** The Vernal BLM RMP/EIS will review the literature on the effects of grazing livestock in arid environments. This review should address the effects of different levels of grazing intensity (stocking rate), grazing systems, range improvements and livestock exclusion on upland habitats. It should include an evaluation of their effects on maintenance, productivity and recovery rates of native plants and soil communities. It should also review the use of various standards, indicators and monitoring methods used to manage grazing and their ability to accurately and timely assess and be representative of plant and soil community characteristics.

**3.2 Determine Soil and Plant Community Status.** Using GIS, map all vegetation, range site types and soil types on BLM lands in the Vernal Resource Area. Design and carry out a science-based review of existing information and a field survey to accurately assess current conditions in the plant and soil community. This survey should determine the distribution of plant species, their ground cover and production. It should do this by allotment, pasture and range site (soil) type. Survey locations should be selected to incorporate distance to the nearest source of water so effects due to proximity to water can be determined. The survey should also include considerations of slope in order to assess the interaction of susceptibility to erosion based on slope and proximity to water sources. Results should be analyzed in the context of past grazing management including type of livestock, seasons of use and grazing intensity. Summarize historical actual use records and grazing systems for each allotment to provide a basis for evaluating proposed actions.

**3.3 Determine Capable and Suitable Acres for Livestock Grazing.** Based on the science we have provided and criteria we have discussed in the documents cited, (slope, distance to water, forage production, hazard of erosion, etc) establish a protocol for determining capable and suitable lands for livestock grazing. This protocol should include a recovery prescription for lands that are severely depleted of their native herbaceous plant communities or have bare ground exceeding thresholds that lead to accelerated erosion.

**3.3 Determine a Sustainable Livestock Stocking Rate.** Establish the percent of diets of domestic sheep, cattle, deer, elk, pronghorn and sage grouse that are grasses, forbs and shrubs and their seasonal variation in selection of these foods. We suggest information from Holechek et al (2001) as a starting point. Using this information and the population management goals for wildlife, calculate wildlife needs. This should include the stubble height and other criteria established in Braun et al (1977) and Connelly et al (2000). Based on the information we provided in section 2.0 above, and the limitations on use of perennial grasses established by the scientific review, calculate the available AUMs of palatable native forage grasses, forbs and shrubs that can be utilized by livestock without impairment of plant productivity. Use this analysis to determine livestock stocking rates for each allotment and pasture for normal and dry precipitation years. Livestock numbers should not be increased during above normal precipitation years to allow for improvement of plant and soil conditions.

**3.4 Develop a Systematic Monitoring Protocol.** Based on the science we have referenced here, develop a systematic monitoring protocol for assessing livestock use (forage amount) in a timely manner to avoid over use of any pasture. The timing of measurement must be established relative to the permitted grazing season of use. It is critical that during the early years of application, these must be assessed frequently in order to ascertain the length of time livestock may remain in each pasture before excessive use occurs. This information will lead to establishment of realistic grazing schedules that lower the risk of excessive livestock grazing and damage. If indicators such as stubble height are proposed, they must also have a demonstrated relationship to actual percent use on grasses, forbs and shrubs by field testing.

**3.5 Determine Recovery Prescriptions for Lands Below Potential.** Analyze peer-reviewed science and government publications we have referenced as well as others that have specifically evaluated different grazing scenarios and their ability to recover native components of damaged arid lands to their potential. Use this information to develop recovery prescriptions for allotments and pastures that are not at potential (based on their departure from potential species distribution, ground cover and productivity). Prescriptions that should be evaluated should include options such as (a) long-term rest, (b) reduced stocking rates, (c) grazing systems, (d) reseeding, (e) vegetation treatments and others. The time required for recovery of potential should be evaluated for each method.

**3.6 Determine Impacts of Water Developments.** Water developments have been used for decades on the promise that they would result in better livestock distribution and improvement in conditions on the ground. BLM has never quantitatively assessed these claims. It is critical that BLM provide this analysis based on evidence from the literature and its own survey of conditions in the Vernal Resource Area. The analysis described in sections **3.1** and **3.2** above should include an element that locates and maps all water developments in the Resource Area on BLM lands. Based on field surveys combined with historical range data and satellite imagery, determine the impacts and/or benefits of these past water developments on upland plant communities and soils. The effects of these water developments on their source waters (spring, stream or seep) and associated wetland habitats should be assessed, documented and reported in the RMP/EIS.

#### **4.0 Management Recommendations for Riparian Areas**

BLM must provide assured protection of springs, streams, riparian areas and wetlands as these areas are highest in wildlife values, yet occupy only a small land area. While continuing to propose additional water developments and grazing systems as means of protecting riparian areas, BLM does not provide scientific evidence that the values of these areas are actually protected by these methods. The evidence in the government and scientific literature is that excluding livestock from riparian areas is the only effective method of protection and recovery.

Don Duff (1977) published results of studies that showed greater fish and vegetation production and stream bank recovery occurred within exclosures on Big Creek, Utah than outside in grazed areas. In fact, he stated that conditions outside the exclosures continued to decline. He concluded that 6 to >8 years of rest were necessary to restore aquatic and riparian habitat. Even after 30 years of excluding livestock, sediment from uplands and riparian areas outside the exclosures continues to impair salmonid spawning, rearing and feeding substrates. The literature is consistent in showing that exclusion of livestock from riparian areas is the most effective method of restoring their ecological integrity. There is little evidence that continued livestock grazing of degraded streams under various management schemes leads to recovery of habitat attributes required for native fish and wildlife. BLM must objectively evaluate its alternatives to livestock exclosures in the literature review proposed in paragraphs (4.1) and (4.7) below and include those evaluations and the science upon which they are based in its RMP/EIS.

We are concerned over the lack of quantitative monitoring of streams, springs and riparian areas. Some of the limitations of the current BLM PFC assessment technique are discussed in Stevens et al (2002). BLM has not provided any systematic protocol for this monitoring and in general does not assess spring or wetland conditions. We are concerned that reliance will be placed on stubble heights without providing any evidence that, in arid ecosystems, use of this standard promotes recovery of critical habitat elements in damaged stream or wetland systems, or for that matter, prevents degradation of those attributes.

If stubble height is used, it must be correlated with percent riparian vegetation use and a stubble height standard set so that livestock use is kept within constraints that do not lead to lowered production of riparian vegetation in future years. In addition, it must be shown to be protective of stream bank integrity through comparison to percent bank trampling, bank erosion and bank stability. The literature review we propose in section **4.1** should address these issues, establish and justify riparian utilization standards and bank trampling standards that are protective of stream banks and riparian vegetation. This analysis of this and other issue presented below must be incorporated into the Vernal BLM RMP/EIS.

**4.1 Literature Review.** The RMP/EIS will review the literature on the effects of livestock grazing, grazing systems, range improvements and livestock exclusion on stream, spring and riparian habitats, their maintenance and recovery as well as water quality. Carter (2001) reviews the effects of livestock grazing on riparian areas and water quality. This report is available on line at:

[http://www.westernwatersheds.org/reports/grazeWQ\\_JCarter/WQWWP.doc](http://www.westernwatersheds.org/reports/grazeWQ_JCarter/WQWWP.doc)

The RMP/EIS will also review the use of various indicators such as riparian stubble height and bank trampling measures, their application and relation to stream bank stability, riparian vegetation productivity, in-stream aquatic habitat and restoration rates. This review will incorporate analysis of the literature we have referenced in addition to any sources BLM and its consultant may provide.

**4.2 Assessment of Current Condition of Springs, Seeps and Wetlands.** Locate and map all springs and wetlands not associated with perennial streams on BLM lands in the Vernal Resource Area. Photograph and describe each, document whether and how they have been modified for water developments and if so, whether the development is still functioning. Assess their current condition, wetland extent (area) and flow using photographs and DOI (1994). These habitats should be surveyed to establish a current baseline. The surveys should also be repeated during each permit mid-term and one year prior to permit renewal. Include the baseline mapping, analysis and photographs in the RMP/EIS.

**4.3 Develop a Monitoring Protocol for Springs, Seeps and Wetlands.** Based on the science reviewed in section **4.1**, develop a systematic monitoring protocol for livestock use of riparian/wetland herbaceous vegetation at springs, streams and wetlands. This assessment protocol will:

- a Establish stubble height and livestock trampling standards and assess these effects at each spring and wetland area and at multiple locations on each

flowing stream within each pasture. A defined protocol should specify the number of measurements to be taken at each location, the length of stream reach to assess and a method to evaluate trampling damage to springs and wetlands.

- b Measure percent use of riparian grasses, not sedges. These measurements will be taken in the riparian zone outside the stream channel, not along the greenline.
- c Monitor water quality for conformance to standards
- d Establish a schedule for each allotment/pasture that measures these parameters no later than 3 weeks after livestock enter a particular pasture or allotment subdivision. It is important to measure these use levels early in the grazing period so stocking rates and management can be adjusted in a timely manner to prevent exceeding the standards. Accelerated measurements during the first year, in particular, are necessary to refine the schedule for future years.
- e Record the data, including allotment, pasture, location designation, turn-in date, monitoring date, riparian species measured, percent bank trampling, percent area trampled (springs and wetlands) and stubble heights. Record raw data on prepared forms so that means and standard deviations can be calculated. Enter this information into an electronic database that can be regularly updated following each monitoring event and is accessible to interested parties either on-line or by request.

**4.4 Assessment of Current Condition of Streams.** Based on best available science, describe a methodology and schedule for performing stream habitat and water quality surveys. This should include analysis of existing data and collection of baseline habitat and water quality data, maps and photographs that will be included in the Vernal RMP/EIS.

**4.5 Develop Stream and Water Quality Monitoring Protocol.** In addition to the baseline survey, the RMP/EIS should develop a monitoring protocol that provides for assessment of stream habitat and water quality during succeeding permit terms near the permit or lease mid-term and the year prior to renewal. This monitoring protocol should continue to include PFC assessments using DOI (1993) as modified by Stevens et al (2002). Winward (2000) also provides insight into stream and riparian monitoring technology. Any of these protocols must be supplemented by collecting additional data that quantitatively documents certain critical stream habitat conditions. We suggest the Rosgen method, USDA (1992, 1997) or other acceptable fish habitat survey protocols. Regardless of method, certain critical pieces of information must be collected to supplement PFC assessments. The literature should be reviewed and analyzed in the RMP/EIS and details of methodologies, data management, reporting and public access explained. This critical supplemental data includes:

- a. Establishing permanent, marked stream cross-section survey points for determining channel profile, width and depth similar to those used by Duff (1977). Priority locations are in Rosgen Type “C” channels, if available and should encompass a complete meander with multiple cross-sections (minimum 3 at each reach). These channel surveys and

- the following data must be collected for a minimum of two reaches of each perennial stream in each pasture.
- b. Bank condition surveys including measurement of the following parameters over a minimum 100' stream reach or full meander on both sides of the stream (Winward, 2000, recommends 363 feet). Bank condition parameters (i – iv\_ will be assessed from the water line to the top of the bank, not at the greenline:
    - i. the linear feet of stream banks that are vegetated/stable
    - ii. the linear feet of stream banks that are vegetated/unstable
    - iii. the linear feet of stream banks that are unvegetated/stable
    - iv. the linear feet of stream banks that are unvegetated/unstable
    - v. the linear feet of stream banks that are undercut
    - vi. the linear feet of stream bank with overhanging vegetation.
  - c. During the baseline survey, and during succeeding permit terms, at permit mid-term and last permit year, collect triplicate MacNeil Core Sediment samples of salmonid spawning gravels (<2") near the downstream extent of each stream in each pasture. Based on the literature, establish criteria for sediment fines (<6.35 mm) as a percent of substrate that allows successful salmonid reproduction. Generally, levels of sediment fines above 30% result in significant mortality of eggs and larvae.
  - d. Data collected will be systematic and quantitative in order that statistical analysis can be employed. All data will be entered into an electronic database for access by interested parties.

In the RMP/EIS, describe the monitoring protocols. This description will include: the actual methodologies, data entry forms, schedules and locations; provide maps of all locations for monitoring; and incorporate an electronic database with links to maps that is accessible to the interested public. Include the baseline data in the RMP/EIS. BLM must show it has the capability, either with its own staff or outside contractors, to do the necessary monitoring to ensure recovery and prevention of future degradation.

**4.6 Develop Stream, Spring and Wetland Recovery Prescriptions.** For streams, springs and wetlands that are not functioning properly and/or have impaired water quality, analyze available methods for their recovery. Include an anticipated time frame for recovery to take place based on the scientific literature. Discuss and evaluate these options for recovery in the RMP/EIS, including their costs and benefits and review the scientific literature provided. Methods for recovery should include consideration of: (a) closing pastures; (b) enclosure fencing; (c) removal of water developments; (d) restoration/replanting of riparian areas; (e) active herding to minimize livestock use; (f) removal of livestock when utilization standards are reached and any other methods discovered in the literature. In the RMP/EIS propose and adopt methods that will best recover these degraded streams during the first permit term. In the on-going permitting, management and monitoring of livestock grazing, the land use plan should require documentation of condition and recovery rates by implementation of the described monitoring program. These will be analyzed to demonstrate the changes resulting from application of the particular prescription or combinations of methods adopted.

Yours truly,



John Carter  
Utah Director

Cc: Planning, Vernal BLM

## **5.0 References**

All references listed are incorporated into our comments and should be used by BLM in preparation of its RMP/EIS.

Anderson, David C., Kimball T. Harper and Ralph C. Holmgren. 1982. Factors influencing development of cryptogamic soil crusts in Utah deserts. *Journal of Range Management* 35(2):180-185.

Anderson, Jay E. and Karl L. Holte. 1981. Vegetation development over 25 years without grazing on sagebrush-dominated rangelands in southeastern Idaho. *Journal of Range Management* 34(1):25-29

Anderson, Jay E. and Richard S. Inouye. 2001. Landscape-scale changes in plant species abundance and biodiversity of a sagebrush steppe over 45 years. *Ecological Monographs* 71(4):531-556.

Armour, C.L., D.A. Duff and W. Elmore. 1991. The effects of livestock grazing on riparian and stream ecosystems. *Fisheries* 16(1): 7 – 11.

Austin, Miriam. 2003a. Water developments and wildlife. Violating the public trust. Red Willow Research.

Austin, Miriam. 2003b. BLM Pleasantview allotment wildlife deaths, 2003. Red Willow Research.

Bartos, Dale L. and Robert B. Campbell, Jr. 1998a. Water depletion and other ecosystem values forfeited when conifer forests displace aspen communities. *Rangeland Management and Water Resources of American Water Works Association*. May 1998: 427-433.

Bartos, Dale L. and Robert B. Campbell, Jr. 1998b. Decline of Quaking Aspen in the Interior West – Examples from Utah. *Rangelands* 20(1):17-24.

Beck, Jeffrey, L. and Dean L. Mitchell. 2000. Influences of livestock grazing on sage grouse habitat. *Wildlife Society Bulletin* 28(4):993-1002.

Belnap, Jayne. 1995. Surface disturbances: their role in accelerating desertification. *Environmental Monitoring and Assessment* 37:39-57.

Belnap, Jayne. 1996. Soil surface disturbances in cold deserts: effects on nitrogenase activity in cyanobacterial-lichen soil crusts. *Biology and Fertility of Soils* 23:362-367.

Belnap, Jayne and D. A. Gillette. 1997. Disturbance of biological soil crusts: impacts on potential wind erodibility of sandy desert soils in southeastern Utah. *Land Degradation and Development* 8:355-362.

Belnap, Jayne, Roger Rosentreter, Steve Leonard, Julie H. Kaltnecker, John Williams and David Eldridge. 2001. Biological soil crusts: ecology and management. U.S. D.O.I. Bureau of Land Management Technical Reference 1730-2.

Belsky, A.J., A. Matzke, and S. Uselman. 1999. Survey of livestock influences on stream and riparian ecosystems in the western United States. *Journal of Soil and Water Conservation* 54(1): 419-431.

Belsky, A. Joy and Dana M. Blumenthal. 1997. Effects of livestock grazing on stand dynamics and soils in upland forests of the Interior West. *Conservation Biology* 11(2):315-327.

Belsky, A. Joy, and Jonathan L. Gelbard. 2000. *Livestock Grazing and Weed Invasions in the Arid West*. A Scientific Report Published by the Oregon Natural Desert Association. Portland, Oregon.

Beymer, Renee J. and Jeffrey M. Klopatek. 1992. Effects of grazing on cryptogamic crusts in pinyon-juniper woodlands in Grand Canyon National Park. *American Midland Naturalist* 127:139-148.

Blaisdell, James P. and Ralph C. Holmgren. 1984. *Managing Intermountain Rangelands – Salt-Desert Shrub Ranges*. USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah. General Technical Report INT-163. 52p.

Blackwell, Jack. 2001. Letter on capability and suitability. Regional Forester, Intermountain Region.

BLM. 1994. *Rangeland Reform '94 Draft Environmental Impact Statement*.

BLM. 2003. *Proposed Revisions to Grazing Regulations for the Public Lands, Draft Environmental Impact Statement DES 03-62*.

Bock, Carl E., Jane H. Bock, William R. Kenney and Vernon M. Hawthorne. 1984. Responses of birds, rodents and vegetation to livestock enclosure in a semidesert grassland site. *Journal of Range Management* 37(3):239-242.

Brotherson, Jack D., Samuel R. Rushforth, and Jeffrey R. Johansen. 1983. Effects of long term grazing on cryptogam crust cover in Navajo National Monument, Ariz. *Journal of Range Management* 36(5):579-581.

Braun, Clait E. 1977. Guidelines for maintenance of sage grouse habitat. *Wildlife Society Bulletin* 5(3):99-105.

Brotherson, Jack D., Samuel R. Rushforth and Jeffrey R. Johansen. 1983. Effects of long-term grazing on cryptogam crust cover in Navajo National Monument, Ariz. *Journal of Range Management* 36(5):579-581.

Bryant, H.T., R.E. Blaser, and J.R. Peterson. 1972. Effect of trampling by cattle on bluegrass yield and soil compaction of a Meadowville loam. *Agronomy Journal*. 64:331-334.

Carlson, Cathy and John Horning. 1992. Big profits at a big price – Public Land ranchers profit at the expense of the range. Publication No. 79950, National Wildlife Federation, 1400 Sixteenth St. N.W. Washington, D.C. 64p.

Carter, John. 1998. Investigation of Spawn Creek, Utah Coliform Contamination and Stream Bank Stability in Relation to Cattle Grazing. Willow Creek Ecology report.

Carter, John G. 2001. Livestock and water quality. Western Watersheds Project Report.

Catlin, James, Joro Walker, Allison Jone, John Carter and Joe Feller. 2003. Multiple use grazing management in the Grand Staircase National Monument. A tool provided to the Monument range staff by the Southern Utah Land Restoration Project.

Clary Warren and Bert Webster. 1989. Managing grazing of riparian areas in the Intermountain Region. Intermountain Research Station, Forest Service. General Technical Report INT-263

Cummins, K.W. and George L. Spengler. 1977. Stream ecosystems. *Water Spectrum*. 10:1-9.

Cummins, K. W. 1974. Stream ecosystem structure and function. *Bioscience* 24:631-641.

Connelly, John W., Michael A. Schroeder, Alan R. Sands and Clait E. Braun. 2000. Guidelines to manage sage grouse populations and their habitats. *Wildlife Society Bulletin* 28(4):967-985.

Cottam, Walter P. 1947. Is Utah Sahara bound? *Bulletin of the University of Utah* 37(11):1-41.

Crider, Franklin J. 1955. Root-growth stoppage resulting from defoliation of grass. Technical Bulletin No. 1102. USDA Soil Conservation Service. 23p

D'Antonio, C. M. and P. M. Vitousek. 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics* 23:63-87.

- Danvir, Rick. Ca2003 (undated). Sage grouse ecology and management in northern Utah sagebrush-steppe. Deseret Land and Livestock, Woodruff, Utah.
- Dodge, Marvin. 1972. Forest fuel accumulation – a growing problem. *Science* 177:139-142.
- DOI. 1993. Riparian Area Management: Process for Assessing Proper Functioning Condition. Department of Interior, BLM TR 1737-9.
- DOI. 1994. Riparian Area Management: Process for Assessing Proper Functioning Condition for Lentic Riparian-Wetland Areas. Department of Interior, BLM TR 1737-11.
- Donahue, Debra L. 1999. The western range revisited – removing livestock from public lands to conserve native biodiversity. University of Oklahoma Press, Norman.
- Duff, D. A. 1977. Livestock grazing impacts on aquatic habitat in Big Creek, Utah. In: Proceedings of the Workshop on Livestock and Wildlife-Fisheries Relationships in the Great Basin. University of California Agric. Station, Sci. Spec. Publication 3301. Berkeley, California.
- Duff, Donald A. 1979. Riparian habitat recovery on Big Creek, Rich County, Utah. In Proceedings: Forum – Grazing and Riparian/Stream Ecosystems. Trout Unlimited, Inc. 91 p.
- Dyksterhuis, E. J. 1949. Condition and management of range land based on quantitative ecology. *Journal of Range Management* 2:104-115.
- Ecosystem Research Institute. 1984. 1983 Annual Report Ecosystem Analysis. Report to White River Shale Oil Corporation, Salt Lake City, Utah. 106p. Report available thru Western Watersheds Project Utah Office at P.O. Box 280, Mendon, Utah 84325.
- EPA. 1998b. The quality of our Nation's water: 1996 – Executive Summary of the National Water Quality Inventory: 1996 Report to Congress. EPA841-S-97-001 Office of Water, U.S. Environmental Protection Agency, Washington, D.C.
- EPA. 1976. Quality criteria for water, July 1976. Fecal coliform bacteria. U.S. Environmental Protection Agency, Washington, D.C.: 42-50
- Evanko, Anthony B. and Roald A. Peterson. 1955. Comparisons of protected and grazed mountain rangelands in southwestern Arizona. *Ecology* 36(1):71-82.
- Fleischner, Thomas L. 1994. Ecological costs of livestock grazing in western North America. *Conservation Biology* 8(3):629-644.
- GAO. 1988. Rangeland management: more emphasis needed on declining and overstocked grazing allotments. U.S. General Accounting Office, GAO/RCED-88-80, Washington, D.C.

- GAO. 1991. Public land management: attention to wildlife is limited. GAO/RCED-91-64
- GAO. 1995. Animal waste management and water quality issues. Publication no. GAO/RCED-95-200BR. 97pp. General Accounting Office, Washington, D.C.
- Galt, Dee, Greg Mendez, Jerry Holechek and Jamus Joseph. 1999. Heavy winter grazing reduces forage production: an observation. *Rangelands* 21(4):18-21
- Galt, Dee, Francisco Molinar, Joe Navarro, Jamus Joseph and Jerry Holechek. 2000. Grazing capacity and stocking rate. *Rangelands* 22(6):7-11.
- Green, D. M. and J. B. Kauffman. 1995. Succession and livestock grazing in a northeastern Oregon riparian ecosystem. *Journal of Range Management* 48:307-313.
- Gregory, Stanley V., Frederick J. Swanson, W. Arthur McKee and Kenneth W. Cummins. 1991. An ecosystem perspective of riparian zones. *Bioscience* 41(8): 540-550.
- Hanley, T. H. and J. L. Page. 1981. Differential effects of livestock use on habitat structure and rodent populations. *California Fish and Game* 68:160-173.
- Harper, K. T., R. Van Buren, and S. Kitchen. 1996. Invasion of alien annuals and ecological consequences in salt desert shrublands of western Utah. Pages 58 -65 in J. R. Barrow, E. Durant, R.E. Sosebee and R. J. Tausch, editors. *Proceedings: Shrubland ecosystem dynamics in a changing environment*. General Technical Report-338. U.S. Forest Service Intermountain Research Station, Ogden, Utah.
- Hild, A.L., M.G. Karl, M.R. Heferkamp and R.K. Heitschmidt. 2001. Drought and grazing III: root dynamics and germinable seed bank. *Journal of Range Management* 54(3):292-298.
- Hobbs, R. J. and L. F. Huenneke. 1992. Disturbance, diversity, and invasion: implications for conservation. *Conservation Biology* 6:324-337.
- Hockett, Glenn A. 2002. Livestock impacts on the herbaceous components of sage grouse habitat: a review. *Intermountain Journal of Sciences* Vol 8(2):105-114.
- Hoffman, L., and R.E. Ries. 1991. Relationship of soil and plant characteristics to erosion and runoff on pasture and range. *Journal of Soil and Water Conservation*, p. 143-147.
- Holechek, Jerry L. 1996a. Financial returns and range condition on southern New Mexico ranches. *Rangelands* 18(2):52-56
- Holechek, Jerry L. 1996b. Drought and low cattle prices: hardship for New Mexico ranchers. *Rangelands* 18(1):11-13.
- Holechek, Jerry L. and Karl Hess, Jr. 1996c. Grazing lands: prices, value, and the future. *Rangelands* 18(3):102-105

- Holechek, Jerry L., Hilton de Souza Gomes, Francisco Molinar and Dee Galt. 1998. Grazing intensity: critique and approach. *Rangelands* 20(5):15-18
- Holechek, Jerry L., Hilton Gomez, Francisco Molinar and Dee Galt. 1999a. Grazing studies: what we've learned. *Rangelands* 21(2):12-16
- Holechek, Jerry L., Milton Thomas, Francisco Molinar and Dee Galt. 1999b. Stocking desert rangelands: what we've learned. *Rangelands* 21(6):8-12
- Holechek, Jerry L., Hilton Gomez, Francisco Molinar, Dee Galt and Raul Valdez. 2000. Short-duration grazing: The facts in 1999. *Rangelands* 22(1):18-22.
- Holechek, Jerry L., Rex D. Pieper and Carlton H. Herbel. 2001. *Range Management: Principles and Practices*, Fourth Edition. Prentice-Hall, New Jersey. 587p
- Howard, Gary L., Steven R. Johnson and Stanley L. Ponce. 1983. Cattle grazing impact on surface water quality in a Colorado front range stream. *J. Soil and Water Conservation*. March-April 1983:124-128.
- Hubbard, R.K., D.L. Thomas, R.A. Leonard and J.L. Butler. 1987. Surface runoff and shallow ground water quality as affected by center pivot applied dairy cattle wastes. *Trans. ASAE* 30(2):430-437.
- Hutchings, S.S. and G. Stewart. 1953. Increasing forage yields and sheep production on Intermountain winter ranges. U.S. Department of Agriculture Circular 925. 63p.
- Jacobs, Lynn. 1991. *Waste of the West*.
- Johansen, Jeffrey R. 1993. Cryptogamic crusts of semiarid and arid lands of North America. *Journal of Phycology* 29:140-147.
- Jones, Allison. 2000. Effects of cattle grazing on North American arid ecosystems: a quantitative review. *Western North American Naturalist* 60(2):155-163.
- Jones, Allison L and William S. Longland. 1999. Effects of cattle grazing on salt desert rodent communities. *The American Midland Naturalist*. 141(1):1-11.
- Julander, Odell. 1962. Range management in relation to mule deer habitat and herd productivity in Utah. *Journal of Range Management* 15(5):278-281.
- Kauffman, J. Boone and W.C. Kreuger. 1984. Livestock impacts on riparian ecosystems and streamside management implications – a review. *Journal of Range Management* 37(5): 430 – 437.
- Kay, Charles E. and Dale L. Bartos. 2000. Ungulate herbivory on Utah aspen: assessment of long-term exclosures. *Journal of Range Management* 53(2):145-153.

- Kay, Charles E. 2001. The condition and trend of aspen communities on BLM administered lands in central Nevada – with recommendations for management. Final Report to Battle Mountain Field Office, Bureau of Land Management.
- Kie, John G. 1996. The effects of cattle grazing on optimal foraging in mule deer (*Odocoileus hemionus*). *Forest Ecology and Management* 88:131-138.
- Kleiner, Edgar F. and K. T. Harper. 1972. Environment and community organization in grasslands of Canyonlands National Park. *Ecology* 53(2):299-309.
- Knick, Steven T., David S. Dobkin, John T. Rotenberry, Michael A. Shroeder, W. Matthew Vander Haegen and Charles Van Riper III. 2003. Teetering on the edge or too late? Conservation and research issues for avifauna of sagebrush habitats. *The Condor* 105:611-634.
- Knopf, Fritz L., James A. Sedgwick and Richard W. Cannon. 1988. Guild structure of a riparian avifauna relative to seasonal cattle grazing. *Journal of Wildlife Management* 52(2):280-290.
- Kreuger, William C. and A. H. Winward. 1974. Influence of cattle and big-game grazing on understory structure of a Douglas-fir Ponderosa Pine- Kentucky bluegrass community. *Journal of Range Management* 27(6):450-453.
- Kreuper, David, Jonathan Bart and Terrell D. Rich. 2003. Response of vegetation and breeding birds to the removal of cattle on the San Pedro River, Arizona (USA). *Conservation Biology* 17(2):607-615.
- Lacey, J. R. 1987. The influence of livestock grazing on weed establishment and spread. *Proceedings: Montana Academy of Science* 47:131-146.
- Lile, David F., Kenneth W. Tate, Donald L. Lancaster and Betsy M. Karle. 2003. Stubble height standards for Sierra Nevada meadows can be difficult to meet. *California Agriculture*, 57(2):60-64.
- Loft, Eric R., John W. Menke and John G. Kie. 1991. Habitat shifts by mule deer: the influence of cattle grazing. *Journal of Wildlife Management*. 55(1):16-26.
- Loope, Walter L., and Neil E. West. 1979. Vegetation in relation to environments of Canyonlands National Park. In: R. M. Linn (ed.) *Proceedings of the First Conference on Scientific Research in the National Parks*. Vol. 1. U.S. Dept. of Interior, NPS Trans. And Proc. Series No. 5.
- Lusby, Gregg C. 1979. Effects of grazing on runoff and sediment yield from desert rangeland at Badger Wash in western Colorado, 1953-73. U.S. Geological Survey Water Supply Paper 1532-1.
- Lusby, Gregg C. 1970. Hydrologic and biologic effects of grazing vs. non-grazing near Grand Junction, Colorado. *Journal of Range Management* 23(4):256-260.

- Mack, R. N. 1981. Invasion of *Bromus tectorum* L. into western North America: An ecological chronicle. *Agro-ecosystems* 7:145-165.
- Mack, R. N. and J. N. Thompson. 1982. Evolution in steppe with few large, hooved mammals. *American Naturalist* 119:757-773.
- Madany, Michael H. and Neil E. West. 1983. Livestock grazing – fire regime interactions within montane forests of Zion National Park, Utah. *Ecology* 64(4):661-667.
- Marble, James R. and Kimball T. Harper. 1989. Effect of timing of grazing on soil surface cryptogamic communities in a great basin low shrub desert: a preliminary report. *Great Basin Naturalist* 49(1):104-107.
- Marcuson, Patrick E. 1977. Overgrazed streambanks depress fishery production in Rock Creek, Montana. Fish and Game Federation Aid Program. F-20-R-21-11a.
- Martin, John H. Jr. 1997. The Clean Water Act and animal agriculture. *J. Environmental Quality* 26:1198-1203.
- McKee, T.B., N.J. Doesken, and J. Kleist, 1993. The relationship of drought frequency and duration of time scales. Eighth Conference on Applied Climatology, American Meteorological Society, Jan 17-23, 1993, Anaheim CA, pp. 179-186.
- McLean, A. and E.W. Tisdale. 1972. Recovery rate of depleted range sites under protection from grazing. *Journal of Range Management* 25:178-184.
- Medin, D. E., and W. P. Clary. 1990. Small mammal populations in a grazed and ungrazed riparian habitat in Nevada. Research Paper INT-413. U.S. Forest Service, Intermountain Research Station, Ogden, Utah.
- Moore, Elbert, Eric Janes, Floyd Kinsinger, Kenneth Pitney and John Sainsbury. 1979. Livestock grazing management and water quality protection. U.S. Environmental Protection Agency and the Bureau of Land Management.
- National Research Council. 2002. Riparian Areas: Functions and Strategies for Management, National Research Council, Committee on Riparian Zone Functioning and Strategies for Management, Water Science and Technology Board, Board on Environmental Studies and Toxicology (National Academy Press, 2002).
- NWF. 1997. Comb Wash decision, *National Wildlife Federation v. Bureau of Land Management*, 140 IBLA 85 (Aug. 21, 1997)
- Orr, Howard K. 1975. Recovery from soil compaction on bluegrass range in the Black Hills. *Transactions of the ASAE* 1076-1081.
- Owens, L.B., W.M. Edwards and R.W. Van Keuren. 1996. Sediment losses from a pastured watershed before and after stream fencing. *Journal of Soil and Water Conservation* 51(1):90-94.

- Packer, Paul. 1998. Requirements for watershed protection on western mountain rangelands. Unpublished manuscript. Dr. Packer is retired from the USDA Intermountain Forest and Range Experiment Station, Logan, Utah.
- Paige, Christine and Sharon A. Ritter. 1999. Managing sagebrush habitats for bird communities. Partners in Flight, Western Working Group, Boise, Idaho.
- Pearce, Richard. 1988. Where deer and cattle roam. USDA Forest Service Pacific Southwest Research Station.
- Pell, Alice N. 1997. Manure and microbes: public and animal health problem? J. Dairy Sci. 80:2673-2681.
- Peterjohn, W.T. and D. L. Correll. 1984. Nutrient dynamics in an agricultural watershed: observations of a riparian forest. Ecology 65: 1466-1475
- Platts, W. S. 1991. Livestock Grazing. *In: Influence of Forest and Rangeland Management on Salmonid Fishes and Their Habitats*. American Fisheries Society Special Publication 19:389-423.
- Ratliff, Joe and Charles Keepports. 2000. Surface water analysis and management recommendations for the Carico Lake Allotment. U.S. Dept. of Interior Bureau of Land Management, Battle Mountain Field Office, Nevada.
- Ray, D.E., A.M. Lane, C.B. Roubicek, and R.W. Rice. 2004. Range beef herd growth statistics. In: Arizona Rancher's Management Guide. Arizona Cooperative Extension, College of Agriculture, University of Arizona.
- Rosenzweig, M. L. and J. Winakur. 1969. Population ecology of desert rodent communities: habitats and environmental complexity. Ecology 50:558-572.
- Rummell, Robert S. 1951. Some effects of livestock grazing on Ponderosa pine forest and range in central Washington. Ecology 32(4):594-607.
- Saxon, Keith E., Lloyd F. Elliott, Robert I. Papendick, Michael D. Jawson and David H. Fortier. 1983. Effect of animal grazing on water quality of non-point runoff in the Pacific Northwest. Project Summary, Robert S. Kerr Environmental Research Laboratory, Ada, Oklahoma. EPA-600/S2-83-071. 7p.
- Schepers, J.S., B.L. Hackes and D.D. Francis. 1982. Chemical water quality of runoff from grazing land in Nebraska: II. contributing factors. J. Environ. Qual., Vol 11(3):355-359.
- Schulz, Terri T and Wayne C. Leininger. 1990. Differences in riparian vegetation structure between grazed areas and exclosures. Journal of Range Management 43(4):295-299.
- Sharp, Lee A., Ken Sanders and Neil Rimbey. 1992. Variability of crested wheatgrass production over 35 years. Rangelands 14(3):153-168

- Schwan, H.E., Donald J. Hodges and Clayton N. Weaver. 1949. Influence of grazing and mulch on forage growth. *Journal of Range Management* 2(3):142-148.
- Stevens, Lawrence E., Peter Stacey, Don Duff, Chad Gourley and James C. Catlin. 2002. Riparian ecosystem evaluation: a review and test of BLM's proper functioning condition assessment guidelines.
- Stewart, Kelley M., R. Terry Bowyer, John G. Kie, Norman J. Cimon, and Bruce K. Johnson. 2002. Temporal distributions of elk, mule deer, and cattle: resource partitioning and competitive displacement. *Journal of Mammalogy*. 83(1):229-244.
- Souder, Jon A. ca1997. How Does Livestock Grazing Fit Into the Larger Societal Uses of Wildlands? Methods for Determining Benefits and Their Application to the Kaibab Plateau. College of Ecosystem Science and Management, Northern Arizona University.
- Taylor, Daniel M. 1986. Effects of cattle grazing on passerine birds nesting in riparian habitat. *Journal of Range Management* 39(3):254-258.
- Thelin, Richard and Gerald F. Gifford. 1983. Fecal coliform release patterns from fecal material of cattle. *Journal of Environmental Quality* 12(1):57-63.
- Tiedemeann, A.R., D.A. Higgins, T.M. Quigley, H.R. Sanderson and D.B. Marx. Responses of fecal coliform in streamwater to four grazing strategies. *Journal of Range Management* 40(4):322-329.
- Trimble, Stanley W. and Alexandra C. Mendel. 1995. The cow as a geomorphic agent – a critical review. *Geomorphology* 13:233-253.
- USDA. 1981. America's soil and water: condition and trends. U.S. Department of Agriculture Soil Conservation Service, Washington, D.C. 33p.
- USDA. 1982. Soil Survey of Rich County Utah. USDA Soil Conservation Service, Forest Service and Bureau of Land Management.
- USDA. 1992. Integrated riparian evaluation guide. U.S. Department of Agriculture, Region IV Forest Service, Ogden, Utah.
- USDA. 1993. Monitoring data from the North Rich allotment, Logan Ranger District, Wasatch-Cache National Forest.
- USDA. 1996. Intermountain Regional Assessment: Properly Functioning Condition. USDA Forest Service, Region IV, Ogden, Utah.
- USDA. 1997. R1/R4 (Northern/Intermountain Regions) Fish and Fish Habitat Standard Inventory Procedures Handbook. USDA Forest Service General Technical Report INT-GTR-346.
- UDWR. 1997. Conservation Agreement and Strategy for Bonneville Cutthroat Trout in Utah. Utah Division of Wildlife Resources Publication 97-19.

- USDA. 1998. Draft Sub-regional Assessment of Properly Functioning Condition for Areas Encompassing the National Forests of Northern Utah. USDA Forest Service, Region IV, Ogden, Utah.
- USDA. 2001. Process Paper I: rangeland capability and suitability for sheep and cattle ranges. Caribou National Forest.
- USDA. 2002. Final Environmental Impact Statement for the Curlew National Grassland. Caribou National Forest.
- U.S. Department of Interior, U.S. Fish and Wildlife Service, U.S. Department of Commerce and U. S. Census Bureau. 2002. 2001 National Survey of Fishing, Hunting and Wildlife-Watching Associated Recreation. 170 p.
- VanHaveren, Bruce P., Eric B. Janes and William L. Jackson. 1985. Nonpoint pollution control on public lands. *Journal of Soil and Water Conservation* p.92-94.
- Van Poolen, H.W. and J. R. Lacey. 1979. Herbage response to grazing systems and stocking intensities. *Journal of Range Management* 32:250-253.
- Walker, Larry. 2002. Impact of grazing on 153 species of southwestern birds. <http://rangenet.org/directory/walker/swbirds.html> .
- Wambolt, C.L., K.S. Walhof and M.R. Frisina. 2001. Recovery of big sagebrush communities after burning in south-western Montana. *Journal of Environmental Management* 61:243-252.
- Welch, Bruce L. 2002 (in editing). Big sagebrush: A sea fragmented into lakes, puddles, and ponds. Gen. Tech. Rep. RMRS-GTR-\_\_\_\_. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: \_\_\_\_p. Data on file at the Shrub Sciences Laboratory, 735 N 500 E, Provo, Utah 84606.
- Welch, Bruce L. and Craig Criddle. 2003. Countering misinformation concerning big sagebrush. Forest Service Rocky Mountain Research Station Research Paper RMRS-RP-40.
- West, Neil E. 1981. Nutrient cycling in desert ecosystems. In: *Arid Land Ecosystems: Structure, Functioning and Management*, Volume 2. Cambridge University Press.
- West, Neil E. 1983. Western Intermountain Sagebrush Steppe. In *Temperate Deserts and Semi-deserts*, edited by N. E. West. Elsevier Scientific Publishing, Amsterdam. p351-373.
- White, Clifford A., Charles E. Olmstead and Charles E. Kay. 1998. Aspen, elk and fire in the Rocky Mountain national parks of North America. *Wildlife Society Bulletin* 26(3):449-462.
- White, Richard K., Robert W. VanKeuren, Lloyd B. Owens, William M. Edwards and Robert H. Miller. 1983. Effects of livestock pasturing on non-point surface runoff.

Project Summary, Robert S. Kerr Environmental Research Laboratory, Ada, Oklahoma. EPA-600/S2-83-011. 6p.

White River Shale Oil Corporation. 1984. Progress Report Environmental Programs 1983. White River Shale Oil Corporation, Salt Lake City, Utah. 785p. Report available thru Western Watersheds Project Utah Office at P.O. Box 280, Mendon, Utah 84325

Winward, Alma. 2000. Monitoring the Vegetation Resources in Riparian Areas. USDA Forest Service General Technical Report RMRS-GTR-47

Winder, J.A., C.C. Bailey, M.G. Thomas and J.L. Holechek. 2000. Breed and stocking rate effects on Chihuahuan Desert cattle production. *Journal of Range Management* 53(1):32-38.

Wuerthner, George and Mollie Matteson. 2002. Welfare ranching: the subsidized destruction of the west. Island Press.

Zimmerman, G. Thomas and L.F. Neuenschwander. 1984. Livestock grazing influences on community structure, fire intensity, and fire frequency within the Douglas-fir/ninebark habitat type. *Journal of Range Management* 37(2):104-110.